

On the NSR formulation of String Theory on $AdS_5 \times S^5$

Dimitri Polyakov[†]

*The Abdus Salam International Centre for Theoretical Physics
Strada Costiera, 11
I-34014 Trieste, ITALIA*

Abstract

We discuss the NSR formulation of the superstring action on $AdS_5 \times S^5$ proposed recently by Kallosh and Tseytlin in the Green-Schwarz formalism. We show that the stress-energy tensor corresponding to the NSR action for AdS superstring contains the brane-like terms, corresponding to exotic massless vertex operators (referred to as the branelike vertices)- the 5-form $e^\phi \psi_{m_1} \dots \psi_{m_5}$ and the 3-form $\partial(e^\phi \psi_{m_1} \dots \psi_{m_3})$, multiplied by ∂X_m . The corresponding sigma-model action has the manifest $SO(1, 3) \times SO(6)$ invariance of superstring theory on $AdS_5 \times S^5$. We argue that adding the branelike terms is equivalent to curving the space-time to obtain the $AdS_5 \times S^5$ background. We commence the study of the proposed NSR sigma-model by analyzing the scattering amplitudes involving the branelike vertex operators. The analysis shows quite an unusual momentum dependence of these scattering amplitudes.

Keywords: PACS:04.50.+h; 11.25.Mj.

December 98

[†] polyakov@ictp.trieste.it

Introduction

Recently the classical type IIB Green-Schwarz (GS) [1] superstring action in the maximally supersymmetric $AdS_5 \times S^5$ background [2,3,4] has been constructed by Metsaev and Tseytlin [5] and later simplified in the work by Kallosh and Tseytlin by fixing the κ -symmetry [6]; see also [7,8]. In accordance with the conjecture that relates the dynamics of $N = 4D = 4$ super Yang-Mills theory with the type IIB superstring theory and supergravity [9,10,11,12,13] in the AdS background, string theory on the AdS space appears to be a plausible candidate for the string-theoretic description of the large N limit of super Yang-Mills theory. On the condition that the analogue of the Maldacena's conjecture is formulated for the non-supersymmetric case and a suitable mechanism is found for supersymmetry breaking, studying the type IIB string dynamics in the $AdS_5 \times S^5$ may be relevant for the understanding of the problem of confinement from the string-theoretic point of view. Some time ago, in an originally independent development we have noticed the existence of certain new exotic massless states in the spectrum of NSR string theory which apparently contain information about non-perturbative aspects of string dynamics. These open string states are described by BRST-invariant two-form and five-form vertex operators of essentially nonzero ghost numbers. They do not correspond to any known massless excitations in perturbative spectrum of an open string; while precise physical interpretation of these vertex operators is still largely a puzzle, in our previous works [14] we have made a preliminary conjecture (though unfortunately still vague and incomplete at the time) that these excitations (which we referred to as branelike states since the superalgebraic arguments point at their connection with M-brane dynamics) may be relevant to the dynamics of non-Abelian gauge theories. In this paper we shall discuss the connection between these two developments; namely we shall attempt to present the NSR formulation of the Kallosh-Metsaev-Tseytlin type action of a type IIB string on AdS; we shall argue that the NSR version of stress-energy tensor corresponding to the superstring action on $AdS_5 \times S^5$ is related to the one of a sigma-model with the background determined by above mentioned branelike excitations. The paper is organized as follows. In the first section we review the branelike states and give physical arguments for their appearance as a result of the internal normal ordering in Ramond-Ramond vertices; then, by using the open string description of D3-branes, we will perform the GS-NSR transformation for the GS stress-energy tensor in the presence of D3-branes and observe the appearance of the terms related to the branelike states.

The resulting NSR stress-energy tensor (as well as the corresponding action) will have the manifest $SO(1,3) \times SO(6)$ symmetry of superstring theory on curved $AdS_5 \times S^5$ background. In the second section we compute and study the properties of scattering amplitudes that involve branelike states. More precisely, we study the influence of the branelike states on the gravitational lensing of the Ramond-Ramond states by D p-branes and observe that at certain values of the momentum the presence of these exotic states leads to additional simple poles in scattering amplitudes.

1. The type IIB action on $AdS_5 \times S^5$ in the NSR formalism and branelike states

Consider the Ramond-Ramond vertex operators at zero momentum in $(-1/2, -1/2)$ and $(-3/2, -3/2)$ -pictures on a disc:

$$\begin{aligned}
V^{(-1/2, -1/2)}(k, z, \bar{z}) &= e^{-\frac{1}{2}\phi}\Sigma^\alpha(z)e^{-\frac{1}{2}\bar{\phi}}\bar{\Sigma}^\beta(\bar{z})e^{ikX}(z, \bar{z})\Gamma_{\alpha\beta}^{m_1\dots m_q}F_{m_1\dots m_q}(k) \\
&\quad V^{(-3/2, -3/2)}(k, z, \bar{z}) \\
&= (e^{-\frac{3}{2}\phi} + \frac{1}{2}e^{\chi-\frac{5}{2}\phi}\partial^2 c)\Sigma^\alpha(z)(e^{-\frac{3}{2}\bar{\phi}} + \frac{1}{2}e^{\bar{\chi}-\frac{5}{2}\bar{\phi}}\bar{\partial}^2 \bar{c}) \\
&\quad \times \bar{\Sigma}^\beta(\bar{z})e^{ikX}(z, \bar{z})\Gamma_{\alpha\beta}^{m_1\dots m_q}F_{m_1\dots m_q}(k)
\end{aligned} \tag{1}$$

where $\phi(z)$ and $\chi(z)$ are free fields that appear in the bosonization of the NSR superconformal ghosts β, γ ; Σ^α is spin operator for NSR matter fields. In the future, unless stated otherwise, we will be dropping the terms depending on fermionic ghosts since these terms will be insignificant for our computations of correlation functions (though in principle these ghost terms are necessary to insure the BRST invariance of vertex operators). If a Ramond-Ramond vertex is placed on a disc and the boundary is present, the holomorphic and anti-holomorphic matter and ghost spin operators are no longer independent but they are related as:

$$\begin{aligned}
\bar{\phi}(\bar{z}) &= \phi(\bar{z}), \bar{\chi}(\bar{z}) = \chi(\bar{z}) \\
\bar{\Sigma}^\alpha(\bar{z}) &= M_\beta^{(p)\alpha}\Sigma^\beta(\bar{z}) \\
M_{\alpha\beta} &\equiv (\Gamma^0 \dots \Gamma^p)_{\alpha\beta}
\end{aligned} \tag{2}$$

The expression for the matrix $M_{\alpha\beta}^{(p)}$ implies that the Dirichlet boundary conditions are imposed on p out of 10 X^m 's while the Neumann conditions are imposed on the rest. As long as the vertices (1) are far from the edge of a D-brane (that is, $z \neq \bar{z}$) one may neglect the interaction between holomorphic and anti-holomorphic spin operators; however, as one approaches the boundary of the disc where $z = \bar{z}$ the internal normal ordering must be performed inside the Ramond-Ramond vertex operators in order to remove the singularities

that arise in the O.P.E. between the spin operators located at z and $\bar{z}$. Adopting the notation $\mathcal{F}^{(q)}(k) \equiv \Gamma^{m_1} \dots \Gamma^{m_q} F_{m_1 \dots m_q}(k)$ we find that the result of the normal ordering is given by:

$$\begin{aligned}
& \lim_{z, \bar{z} \rightarrow s} : e^{-\frac{3}{2}\phi} \Sigma_\alpha : (z) \mathcal{F}_{\alpha\beta}^{(q)} : e^{-\frac{3}{2}\bar{\phi}} \bar{\Sigma}_\beta(\bar{z}) : \\
& \sim \frac{1}{z - \bar{z}} \text{Tr}(\mathcal{F}^{(q)} M^{(p)} \Gamma^{m_1 \dots m_5}) e^{-3\phi} \psi_{m_1} \dots \psi_{m_5}(s) + \dots \\
& \lim_{z, \bar{z} \rightarrow s} : e^{-\frac{3}{2}\phi} \Sigma_\alpha : (z) \mathcal{F}_{\alpha\beta}^{(q)} : e^{-\frac{1}{2}\bar{\phi}} \bar{\Sigma}_\beta(\bar{z}) : \\
& \sim \frac{1}{z - \bar{z}} \text{Tr}(\mathcal{F}^{(q)} M^{(p)} \Gamma^{m_1 m_2}) e^{-2\phi} \psi_{m_1} \psi_{m_2}(s) + \dots
\end{aligned} \tag{3}$$

where we have dropped the less singular terms in the O.P.E. as well as full derivatives. We see that due to the internal normal ordering at the boundary of the disc the Ramond-Ramond vertex operators degenerate into massless *open – string* vertices - the two-form $Z_{mn} = e^{-2\phi} \psi_m \psi_n$ and the five-form $Z_{m_1 \dots m_5} = e^{-3\phi} \psi_{m_1} \dots \psi_{m_5}$. As for the five-form state, there also exists the “dual” version of it in the +1-picture: $Z_{m_1 \dots m_5}^{(+1)} \equiv: \Gamma^4 Z_{m_1 \dots m_5} := e^\phi \psi_{m_1} \dots \psi_{m_5}$ where Γ^4 is the normally ordered fourth power of picture-changing operator. Actually it is this picture-changed version of the 5-form (i.e. the one in the +1-picture) that will appear in the NSR formulation of the AdS superstring action.

As it is the well-known fact that there are no two-form and five-form particles in the *peturbative* spectrum of an open string, giving a proper physical interpretation to these *new* massless states in the spectrum of a superstring is a challenging puzzle. In our previous works [14] we have shown that two-form and five-form vertices (3) appear as central terms in the space-time superalgebra for NSR superstring theory when the supercharges are taken in non-canonical pictures. Since p-form central terms in a SUSY algebra are always related to the presence of p-branes we have argued that the operation of picture-changing (applied to space-time supercharges) is a worldsheet interpretation of the S-duality transformation and the open string vertex operators (3) have essentially non-perturbative origin and represent the M-branes, though the precise relation of these vertex operators to the dynamics of branes is still unclear. In the current paper we shall discuss the role that these new superstring states may play in the recently proposed AdS-CFT correspondence [13,12,9,10,11] and in building the string-theoretic approach to Yang-Mills theories. In the light of the newly found relation between the Yang-Mills correlation functions and the minimum of the supergravity action on the AdS space, the type IIB superstring theory on AdS (which has the AdS supergravity its low-energy limit) appears a plausible candidate for the ansatz to describe the dynamics of large N gauge theories. It

is therefore crucial to understand the dynamics of string theory in the AdS background in order to deepen our understanding of Yang-Mills dynamics and CFT in various dimensions. In this paper we shall claim that in terms of NSR formalism superstring theory on the AdS space (presumably describing the large N limit of gauge theories) corresponds to a sigma model with *flat* space-time metric and with massless background fields corresponding to the branelike vertex operators (2). There are two dual ways of regarding the superstring dynamics in the field of D-branes. The first way is to consider the closed GS superstring propagating in the AdS background corresponding to the near-horizon geometry of D3-brane solution. The $AdS_5 \times S^5$ action in the GS formalism is given by [6]:

$$S = -\frac{1}{2} \int d^2z \{ \sqrt{-g} g^{ij} (y^2 (\partial_i x^p - 2i\bar{\theta} \Gamma^p \partial_i \theta) (\partial_j x^p - 2i\bar{\theta} \Gamma^p \partial_j \theta) + \frac{1}{y^2} \partial_i y^t \partial_j y^t) + 4\epsilon^{ij} \partial_i y^t \bar{\theta} \Gamma^t \partial_j \theta \} \quad (4)$$

where $X^p, p = 0...3$ and $Y^t, t = 4...10$ are the longitudinal and transverse coordinates with respect to the worldvolume of the D3-branes. The second way to explore the GS superstring dynamics in the D-brane background (which is equivalent up to the D-brane's massive modes) is to study the propagation of the GS closed superstring in the *flat* space-time in the presence of open Dirichlet strings [15,16]. In the presence of open strings the holomorphic and antiholomorphic fields θ and $\bar{\theta}$ are no longer independent. The GS superstring action in the flat space-time is given by:

$$-\frac{1}{2} \int d^2z \{ (\partial X^m - i\bar{\theta} \Gamma^m \partial \theta) (\bar{\partial} X_m - i\bar{\theta}^\alpha (\Gamma_{\alpha\beta})_m \bar{\partial} \theta^\beta) - 2i\epsilon^{ij} \partial_i X^m \bar{\theta}^\alpha (\Gamma_{\alpha\beta})_m \partial_j \theta^\beta \} \quad (5)$$

The corresponding stress-energy tensor is given by

$$T_{ij} = \Pi_i^m \Pi_{mj} \quad (6)$$

$$\Pi_i^m = \partial_i X^m + i\bar{\theta}^\alpha \Gamma_{\alpha\beta}^m \partial_i \theta^\beta$$

In the flat space-time one may use the standard relations that exist (up to picture-changing transformation) between GS fermionic variable θ and NSR spin operators in ten-dimensional space-time: $\theta^\alpha = e^{\frac{\phi}{2}} \Sigma^\alpha + \text{ghosts}$.

In the presence of the D3-branes the space-time fermionic fields θ and $\bar{\theta}$ are not independent but are related to each other according to (2). For the D3-branes this relation is given by:

$$\bar{\theta}_\alpha(\bar{z}) = [i(\Gamma^0 \dots \Gamma^3)_{\alpha\beta}] \theta_\beta(\bar{z}) \quad (7)$$

Using (7) let us now proceed with the formulation of the NSR analogue of the stress-energy tensor (6). Because of (7) the normal reordering between $\bar{\theta}$ and θ must be made, similar to the one performed in the Ramond-Ramond vertex operators in (3). Using (7) and the formula for the O.P.E. between two spin operators:

$$: \Sigma_{\alpha}(z) :: \Sigma_{\beta}(w) : \sim \frac{\epsilon_{\alpha\beta}}{(z-w)^{\frac{5}{4}}} + \sum_p \frac{\Gamma_{\alpha\beta}^{m_1 \dots m_p} \psi_{m_1} \dots \psi_{m_p}}{(z-w)^{\frac{5}{4}-p}} \quad (8)$$

one finds that the stress-energy tensor T_{zz} of (6) rewritten in the NSR formalism is given by:

$$\begin{aligned} T_{zzNSR} = & \frac{1}{2} [\partial X_m \partial X^m + \{e^{\phi} (\partial \phi \partial \psi_{\mu} \partial X^{\mu} + \psi_m \partial \psi^m \psi_n \partial X^n) \\ & + \epsilon^{p_1 p_2 p_3 p_4} [e^{\phi} \psi_{p_1} \psi_{p_2} \psi_{p_3} \psi_{p_4} \psi^t \partial y^t + \partial (e^{\phi} \psi_{p_1} \psi_{p_2} \psi_{p_3}) \partial \tilde{x}_{p_4}] \\ & + \frac{1}{2} : \Gamma \Gamma \psi \partial \psi : + ghosts] \end{aligned} \quad (9)$$

Here $\epsilon^{p_1 \dots p_4}$ is the rank 4 antisymmetric tensor for the D3 brane worldvolume (the indices $p_1, \dots, p_4$ run from 0 to 3, t runs from 4 to 9); $m = (p, t)$ and n are ten-dimensional indices, $X^m \equiv (\tilde{x}^p, y^t)$. We have chosen the conformal gauge for the worldsheet metric and dropped full derivative terms. The tensor (9) may be rewritten in the form where the kinetic term for the NSR fermions (quadratic in ψ) appears in a more transparent way:

$$\begin{aligned} T_{zzNSR} = & [\frac{1}{2} \partial X^m \partial X_m + \{\frac{1}{2} : (\Gamma + \Gamma \Gamma) \psi^m \partial \psi_m : \\ & + \epsilon^{p_1 p_2 p_3 p_4} [e^{\phi} \psi_{p_1} \psi_{p_2} \psi_{p_3} \psi_{p_4} \psi^t \partial y^t + \partial (e^{\phi} \psi_{p_1} \psi_{p_2} \psi_{p_3}) \partial \tilde{x}_{p_4}] + ghosts] \end{aligned} \quad (10)$$

Here $\Gamma \equiv: \delta(\gamma)(G_{matter} + G_{ghost}) :$ is the picture-changing operator where G is the matter + ghost supercurrent. Up to the picture-changing transformation, the kinetic part of the resulting NSR stress-energy tensor is the same as in the NSR tensor in flat ten-dimensional space-time without branes. The stress-energy tensor (10) corresponds to a sigma-model with flat space-time metric and the branelike part; it is the branelike states that break the $SO(1,9)$ Lorentz symmetry of the flat space-time action to $SO(1,3) \times SO(6)$ of that on $AdS_5 \times S^5$. The branelike part corresponds to the term $\sim \bar{\theta}^{\alpha} \Gamma_{\alpha\beta}^m \partial \theta^{\beta} \partial X_m$ in the stress-energy tensor. It is important to note that it is crucial to have the D-3 branes (that give rise to the curved $AdS_5 \times S^5$ background in the “dual” approach) in order to include the branelike states (3) in the sigma-model; in the absence of D3-branes (that is, in the flat space-time) there would be no $\Gamma^0 \dots \Gamma^3$ factor in (7) and the 5-form and the 3-form terms in (9), (10) would have vanished. For instance, in the absence of the above gamma-matrix

factor the corresponding 5-form term would be given by $\sim Tr(\Gamma^{\mu_1 \dots \mu_5} \Gamma^\nu) e^\phi \psi_{\mu_1} \dots \psi_{\mu_5} \partial X_\nu$ but this expression obviously vanishes since, due to anticommutation relations between the gamma-matrices, one would inevitably have to contract a pair of anticommuting fermions ψ_μ, ψ_ν with the Minkowski metric tensor $\eta^{\mu\nu}$. In the next section of this paper we will explore the properties of the sigma-model corresponding to the energy-momentum tensor (10) and study the correlation functions of the branelike vertex operators. The calculation will show a rather peculiar momentum dependence of these correlators.

Correlation Functions of Branelike States

Rather than computing the amplitudes involving the open string branelike states (3) in a straightforward way we will choose, for the sake of simplicity, to study the problem of gravitational lensing of the the Ramond-Ramond particles in the non-standard $(-3/2, -3/2)$ -picture in the D-brane field which in some way is equivalent to studying the properties of branelike states (see the comment below). In case of the RR particles in the *standard* $(-1/2, -1/2)$ picture the appropriate scattering amplitude has been computed in [16] The essential difference between these two computations is related to the fact that the Ramond-Ramond vertex $V^{(-3/2, -3/2)}$ of (1) (unlike the RR vertex in the standard picture) evolves into the exotic five-form state (3) on the boundary of the disc, as we have shown above. Therefore one should expect that the scattering amplitude for the RR states in the D-brane field is *picture – dependent*. The computation will show that in comparison with the standard scattering amplitude for the $(-1/2, -1/2)$ RR particles there are additional simple poles that originate from integrating over the region $r \sim 1$ (i.e. near the boundary of the disc) where the $(-3/2, -3/2)$ RR vertex operator (2) becomes the branelike 5-form state (3), therefore these extra poles in the scattering amplitude are apparently of the branelike origin. Consider the correlation function:

$$\begin{aligned}
W(z, \bar{z}, w, \bar{w}, k_1, k_2) &= \langle e^{-\frac{3}{2}\phi} \Sigma_{\alpha_1}(z) e^{-\frac{3}{2}\bar{\phi}} \bar{\Sigma}_{\beta_1}(\bar{z}) F_{\alpha_1 \beta_1}^{(p)} e^{ik_1 X}(z, \bar{z}) \\
&\quad \times e^{\frac{\phi}{2}} \Sigma_{\gamma_2} \Gamma_{\gamma_2 \alpha_2}^m \partial X_m(w) e^{\frac{\bar{\phi}}{2}} \bar{\Sigma}_{\delta_2} \Gamma_{\delta_2 \beta_2}^n \bar{\partial} X_n F_{\alpha_2 \beta_2}^{(p)}(k_2) e^{ik_2 X}(w, \bar{w}) \rangle \\
&= (F^{(p)}(k_1) M^{(q)})_{\alpha_1 \rho_1} (\Gamma^m F^{(p)}(k_2) \Gamma^n M^{(q)})_{\gamma_2 \rho_2} \langle e^{-\frac{3}{2}\phi}(z) e^{-\frac{3}{2}\bar{\phi}}(\bar{z}) e^{\frac{\phi}{2}}(w) e^{\frac{\bar{\phi}}{2}}(\bar{w}) \rangle \\
&\quad \times \langle \Sigma_{\alpha_1}(z) \Sigma_{\rho_1}(\bar{z}) \Sigma_{\gamma_2}(w) \Sigma_{\rho_2}(\bar{w}) \rangle \\
&\quad \times \langle e^{ik_1 X}(z) e^{ik_1 X^\pm}(\bar{z}) \partial X_m e^{ik_2 X}(w) f(n) \partial X_n e^{ik_2 X^\pm}(\bar{w}) \rangle
\end{aligned} \tag{11}$$

Here $f(n) = -1$ for the longitudinal indices (with respect to the D3 brane worldvolume) and 1 for the transverse ones, $X^\pm \equiv f(n)X$. To compute the X -independent part W_0 of this correlation function (that involves ghosts and fermions) one notes that

$$W_0 \sim \frac{\langle e^{-\frac{3}{2}\phi} e^{-\frac{3}{2}\bar{\phi}} e^{\frac{\phi}{2}} e^{\frac{\bar{\phi}}{2}} \rangle}{\langle e^{-\frac{\phi}{2}} e^{-\frac{\bar{\phi}}{2}} e^{-\frac{\phi}{2}} e^{-\frac{\bar{\phi}}{2}} \rangle} \langle \Lambda \Lambda \Lambda \Lambda \rangle \tag{12}$$

where Λ are the full (ghost + matter) (1,0) spin operators in the standard $-\frac{1}{2}$ -picture.

A bit long but elementary computation gives the following result for the zero momentum part of the correlator:

$$W(z, \bar{z}, w, \bar{w}, k_{1,2} = 0) = Tr(\Gamma^r \not{F}^{(p)}(k_1) M^{(q)}) Tr(\Gamma^r \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)}) \frac{|z - \bar{w}|^2}{(w - \bar{w})(z - \bar{z})^3} \\ + 8Tr(\not{F}^{(p)}(k_1) M^{(q)} \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)}) \frac{1}{(z - \bar{z})^2} \quad (13)$$

The kinetic part of the correlator is given by:

$$\langle e^{ik_1 X}(z) e^{ik_1 X^\pm}(\bar{z}) \partial X_m e^{ik_2 X}(w) f(n) \partial X_n e^{ik_2 X^\pm}(\bar{w}) \rangle \\ = \delta(k_1^\parallel + k_2^\parallel) [(z - \bar{z}) \{k_1^2\} (w - \bar{w}) \{k_2^2\} |z - \bar{w}|^2 \{k_1 k_2\} |z - w|^{2(k_1 k_2)}] \\ \times [\frac{\delta_{mn} f(n)}{(w - \bar{w})^2} + (\frac{k_{2n} f(n)}{w - \bar{w}} + k_{1n} f(n) (\frac{1}{z - \bar{w}} - \frac{1}{\bar{z} - w}) + k_{1n} (\frac{1}{z - w} - \frac{1}{\bar{z} - \bar{w}})) \\ \times (\frac{k_{2m} f(m)}{w - \bar{w}} + k_{1m} f(m) (\frac{1}{z - \bar{w}} - \frac{1}{\bar{z} - w}) + k_{1m} (\frac{1}{z - w} - \frac{1}{\bar{z} - \bar{w}})) \\ + \frac{\delta_{mn}}{w - \bar{w}} (2\{k_1 k_2\} (\frac{1}{z - \bar{w}} - \frac{1}{\bar{z} - w}) + 2(k_1 k_2) (\frac{1}{\bar{z} - \bar{w}} - \frac{1}{z - w}))] \quad (14)$$

where we defined $\{k_1, k_2\} \equiv k_{1p} k_2^p - k_{1t} k_2^t \equiv (k_1 k_2)^\parallel - (k_1 k_2)^\perp$ and similarly for $\{k_{1,2}^2\}$; $(k_1 k_2)$ stands for the regular scalar product and $k^\parallel \equiv k_p k^p; p = 0, \dots, q; t = q + 1, \dots, 9$.

So far we were considering the correlation functions on a half-plane. Let us make the conformal transformation $(z, \bar{z}) \rightarrow (u, \bar{u})$ from the half-plane to the disc:

$$z = \frac{i}{2} \left(\frac{u + i}{u - i} \right) \quad (15)$$

Let us furthermore fix $w = -\frac{i}{2}$ so that its image is at the center of the disc. In terms of the disc coordinates $u \equiv r e^{i\varphi}$ the correlation function (14) becomes:

$$W(r, \varphi, k_1, k_2) = (r^2 + 1 - 2r \sin \varphi)^{-\{k_1^2\} - \{k_1 k_2\} - (k_1 k_2)} (r^2 - 1)^{2(k_1^\parallel)^2} r^{2(k_1 k_2)} \\ \times [Tr(\Gamma^r \not{F}(k_1)^{(p)} M^{(q)}) Tr(\Gamma^r \Gamma^m \not{F}(k_2)^{(p)} \Gamma^n M^{(q)}) \frac{(r^2 + 1 - 2r \sin \varphi)^2}{(r^2 - 1)^3} \\ + 8Tr(\not{F}^{(p)}(k_1) M^{(q)} \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)}) \frac{(r^2 + 1 - 2r \sin \varphi)^2}{(r^2 - 1)^2}] \\ \times [\delta_{mn} f(n) + (k_{2n} f(n) + 4k_{1n} r \cos \varphi + k_{1n} (1 + \frac{\sin \varphi}{r})) \\ \times (k_{2m} f(m) + 4k_{1m} r \cos \varphi + k_{1m} (1 + \frac{\sin \varphi}{r})) \\ + \delta_{mn} (8\{k_1 k_2 r \cos \varphi + 2(k_1 k_2) (1 + \frac{\sin \varphi}{r}))] \quad (16)$$

Since the momentum in the longitudinal directions (with respect to the D3-brane world-volume) is conserved one has $\{k_1^2\} + \{k_1 k_2\} + (k_1 k_2) = 0$. Integrating over r and φ we obtain the following expression for the amplitude:

$$\begin{aligned}
A(k_1, k_2) &\equiv \int_0^1 r dr \int_0^{2\pi} d\varphi W(r, \varphi, k_1, k_2) \\
&= 4k_{1n}k_{1m}B(4 + (k_1 k_2), 2(k_1^{\parallel})^2 - 2)Tr(\Gamma^r \not{F}(k_1)^{(p)} M^{(q)})Tr(\Gamma^r \Gamma^m \not{F}(k_2)^{(p)} \Gamma^n M^{(q)}) \\
&\quad + 8Tr(\not{F}^{(p)}(k_1) M^{(q)} \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)})B(4 + (k_1 k_2), 2(k_1^{\parallel})^2 - 1) \\
&\quad + (Tr(\Gamma^r \not{F}(k_1)^{(p)} M^{(q)})Tr(\Gamma^r \Gamma^m \not{F}(k_2)^{(p)} \Gamma^n M^{(q)})B(3 + (k_1 k_2), 2(k_1^{\parallel})^2 - 2) \\
&\quad + 8Tr(\not{F}^{(p)}(k_1) M^{(q)} \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)})B(3 + (k_1 k_2), 2(k_1^{\parallel})^2 - 1)) \\
&\quad \times (5k_{1m}k_{1n} + (2(k_1 k_2) + f(n))\delta_{mn} + k_{2n}k_{2m}f(n)) \\
&\quad + (Tr(\Gamma^r \not{F}(k_1)^{(p)} M^{(q)})Tr(\Gamma^r \Gamma^m \not{F}(k_2)^{(p)} \Gamma^n M^{(q)})B(2 + (k_1 k_2), 2(k_1^{\parallel})^2 - 2) \\
&\quad + 8Tr(\not{F}^{(p)}(k_1) M^{(q)} \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)})B(2 + (k_1 k_2), 2(k_1^{\parallel})^2 - 1)) \\
&\quad \times \left(\frac{7}{2}k_{1m}k_{1n} + (4(k_1 k_2) + 3f(n))\delta_{mn} \right. \\
&\quad \left. + 3k_{2n}k_{2m}f(m)f(n) - k_{1m}k_{2n}f(n) - k_{2m}k_{1n}f(m)\right) \\
&\quad + (Tr(\Gamma^r \not{F}(k_1)^{(p)} M^{(q)})Tr(\Gamma^r \Gamma^m \not{F}(k_2)^{(p)} \Gamma^n M^{(q)})B(1 + (k_1 k_2), 2(k_1^{\parallel})^2 - 2) \\
&\quad + 8Tr(\not{F}^{(p)}(k_1) M^{(q)} \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)})B(1 + (k_1 k_2), 2(k_1^{\parallel})^2 - 1)) \\
&\quad \times \left(\frac{3}{2}k_{1m}k_{1n} + \delta_{mn}f(n) + k_{2m}k_{2n}f(m)f(n) - k_{1n}k_{2m}f(n) - k_{1m}k_{2n}f(n)\right) \\
&\quad + \frac{1}{2}(Tr(\Gamma^r \not{F}(k_1)^{(p)} M^{(q)})Tr(\Gamma^r \Gamma^m \not{F}(k_2)^{(p)} \Gamma^n M^{(q)})B((k_1 k_2), 2(k_1^{\parallel})^2 - 2) \\
&\quad + 8Tr(\not{F}^{(p)}(k_1) M^{(q)} \Gamma^m \not{F}^{(p)}(k_2) \Gamma^n M^{(q)})B((k_1 k_2), 2(k_1^{\parallel})^2 - 1))k_{1m}k_{1n}
\end{aligned} \tag{17}$$

where $B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$ is the Euler beta function. Some comments should be made about the momentum behaviour of the amplitude (17). First of all it is noteworthy that, in addition to the poles present in the “standard” scattering amplitude of [16] there are additional physical singularities (i.e. the simple poles) in the amplitude (17) at $(k_1^{\parallel})^2 = 0, \frac{1}{2}, 1$ that originate from integration over the region $r \sim 1$, i.e. near the boundary of the disc where, as we pointed out earlier, the Ramond-Ramond vertex in the $(-3/2, -3/2)$ -picture becomes the 5-form branelike state (3). One may therefore conclude that it is exclusively the *boundary* 5-form state (3) that gives rise to the additional simple poles in the scattering amplitude (17). Note that these poles did not arise in the calculation in [16] as the RR vertex in the $(-1/2, -1/2)$ -picture degenerates simply into the vector

emission vertex at $r \sim 1$. Therefore the boundary contribution in [16] had no peculiarities. In the $(-3/2, -3/2)$ case the situation is quite different. Note that three additional poles depend exclusively on the momentum of the incoming $(-3/2, -3/2)$ Ramond-Ramond particle (and consequently, on the momentum of the 5-form branelike state). Such an unusual momentum dependence may be the consequence of the fact that the 5-form vertex operator (3) is physical at some discrete values of momentum only (such a behaviour would be reminiscent of that of discrete states in two-dimensional gravity). Another difference with the scattering amplitude in the $(-1/2, -1/2)$ -picture is that in case of the D-instanton background ($q = -1$) the $(-3/2, -3/2)$ scattering amplitude does *not* have the field-theoretic structure observed in [16], on the contrary due to the presence of the 5-form state the amplitude diverges (since $k_1^{\parallel} = 0$ for the D-instanton background). As for the two-form branelike vertex operator $\sim e^{-2\phi} \psi_m \psi_n e^{ikX}$, there is also a peculiarity at $k = 0$. Namely, at any nonzero value of the momentum k the two-form vertex (3) may be decomposed into a sum of a fermionic part of a vector vertex operator in the -2 -picture plus BRST-trivial part:

$$\begin{aligned}
e^{-2\phi} \psi_m \psi_n e^{ikX} &= A_{mn} + B_{mn} \\
A_{mn} &= \frac{1}{k^2} e^{-2\phi} (k\psi)(k_m \psi_n - k_n \psi_m) \\
B_{mn} &= e^{-2\phi} \psi_m \psi_n e^{ikX} - A_{mn}
\end{aligned} \tag{18}$$

To show that one has to consider the correlation function of $e^{-2\phi} \psi_m \psi_n e^{ikX}$ with two other vector emission vertices in the 0-picture on the sphere. It is then straightforward to see that

$$\begin{aligned}
\langle e^{-2\phi} \psi_m \psi_n e^{ikX} V^0(p_1, w_1) V^0(p_2, w_2) \rangle &= \langle A_{mn} V^0(p_1, w_1) V^0(p_2, w_2) \rangle \\
\langle B_{mn} V^0(p_1, w_1) V^0(p_2, w_2) \rangle &= 0
\end{aligned} \tag{19}$$

where $V^0(p, w) = e_m(p)(\partial X^m + i(p\psi)\psi^m)e^{ipX}$. Therefore at a nonzero momentum the two-form vertex operator (3) can be expressed in terms of the picture -2 vector emission vertex, however at $k = 0$ such a decomposition is no longer possible and the two-form becomes a new massless state.

Conclusion

One of the claims of this work is that the superstring action on the $AdS_5 \times S^5$ can be viewed, from the point of view of NSR formalism, as a sigma-model on the *flat* spacetime; introducing the background terms - a 5-form and a 3-form branelike terms is equivalent to curving the $SO(1,9)$ flat space-time to obtain the AdS background. The $SO(1,3) \times SO(6)$ symmetry of superstring theory on $AdS_5 \times S^5$ space is manifest in the energy-momentum tensor (10). Therefore we hope that exploring the dynamics of superstring theory on $AdS_5 \times S^5$ may be analogous to studying the correlation functions of branelike vertex operators on the sphere. As for the action corresponding to the stress-energy tensor (10) one should also add terms corresponding to the Ramond-Ramond background, corresponding to non-vanishing RR scalar field in the near-core D-instanton background [17] These Ramond-Ramond terms should correspond to the WZ term in the action (4) (which does not contribute to the stress- energy tensor). The NSR action corresponding to the stress-energy tensor (10) should, up to yet unknown RR part, look like

$$S_{NSR} = \int d^2z \left[\frac{1}{2} \partial X^m \bar{\partial} X_m + \left\{ \frac{1}{2} : (\Gamma + \Gamma \bar{\Gamma}) \psi^m \bar{\partial} \psi_m : \right. \right. \\ \left. \left. + \epsilon^{p_1 p_2 p_3 p_4} [e^\phi \psi_{p_1} \psi_{p_2} \psi_{p_3} \psi_{p_4} \psi_t \bar{\partial} y^t + \partial(e^\phi \psi_{p_1} \psi_{p_2} \psi_{p_3}) \bar{\partial} \tilde{x}_{p_4}] + ghosts + RR - terms \right] \quad (20)$$

Note that since the 5-form branelike state has vanishing s-matrix elements among elementary string states (see also the discussion in the Appendix), adding branelike terms to the NSR action in flat space-time does not affect perturbative string amplitudes and does not create problems with unitarity. Because of the ghost anomaly cancelation condition adding the 5-form term to the action would not affect the 2-point amplitude of [16] where the RR vertex operators are taken in the standard picture. Another crucial problem is related to the fact that the branelike terms in the NSR stress-energy tensor (10) (as well as possible Ramond-Ramond terms) have essentially nonzero ghost number which cannot be removed by picture-changing transformation. Therefore in order to explore the properties of the sigma-model with the stress-energy tensor (10) it is crucial to introduce some function of picture-changing operators in the measure of functional integration in order to insure the correct ghost number balance (that would cancel the ghost number anomaly). The proper choice of such a measure deformation is an open question. We hope that the conditions of the worldsheet conformal invariance (i.e. vanishing of the 1-loop beta-function) which should be related to requiring the functional with the action corresponding to the energy-momentum tensor (10) to satisfy the large N loop equation, as well as conditions for the amplitude factorization will be sufficient to determine the correct deformation of the measure.

Acknowledgements

It is a pleasure to thank M.Douglas, N.Hambli, A.M.Polyakov and S.Shatashvili for very important suggestions and discussions. The author would like to express his deep gratitude to the European Postdoctoral Institute (EPDI) for the support during the years of 1996-1998 and to the Institut des Hautes Etudes Scientifiques (IHES) for the hospitality during the celebration of the 40th anniversary of the IHES where I elaborated on basic ideas of this work. The author gratefully acknowledges the hospitality and support of the Abdus Salam International Centre for Theoretical Physics (ICTP) in Trieste and especially thanks S.Randjbar-Daemi. I'm also thankful to the physics department of the University of Rome 2 "Tor Vergata" for the hospitality during the academic year of 1997/98 (in particular it is a pleasure to thank M.Bianchi and A.Sagnotti).

Appendix

We would like to comment briefly on the correlation functions of the 5-form branelike state (3) with elementary string states to demonstrate the vanishing of these correlators. The vanishing of these correlation functions allows one to add the branelike terms to the flat NSR action without creating problems with the unitarity. Let us consider first the correlator of the 5-form state with the vector emission vertices on the sphere. The vector emission vertices in various pictures are given by:

$$\begin{aligned} V^{(0)} &= e_m(k)(\partial X^m + i(k\psi)\psi^m)e^{ikX} \\ V^{(-1)}(k) &= e_m(k)e^{-\phi}\psi^me^{ikX} \\ V^{(-2)}(k) &= e^{-2\phi}V^{(0)}(k) \end{aligned} \tag{21}$$

On the sphere the total ghost number of correlators must be equal to -2 to compensate the ghost anomaly. First of all, it is clear that the 3-point function and the 4-point function vanish:

$$\langle V^{(-2)}(k_1)V^{(-1)}(k_2)e^\phi\psi_{m_1}\dots\psi_{m_5}\rangle = \langle V^{(-1)}(k_1)V^{(-1)}(k_2)V^{(-1)}(k_3)e^\phi\psi_{m_1}\dots\psi_{m_5}\rangle \equiv 0 \tag{22}$$

simply because in both cases the total number of worldsheet fermions ψ in the vector vertices (taken together) is less than 5 - so that there are some ψ 's left in the five-form vertex operator with no partner to contract with. The $n > 4$ -point correlation functions vanish also. For the sake of brevity, let us demonstrate it on the example of the 5-point function, for other n -point functions the argument remains the same. Let us introduce the notation $V_5 = e^\phi\psi_{m_1}\dots\psi_{m_5}B^{m_1\dots m_5}$ where B is some rank 5 tensor which, without a loss of generality, may be assumed antisymmetric. The 5-point function is given by:

$$W_5 = \langle V^{(0)}(k_1)V^{(-1)}(k_2)V^{(-1)}(k_3)V^{(-1)}(k_4)V_5 \rangle \tag{23}$$

Clearly, it is only the fermionic part of the vertex $V^{(0)}$ that could, in principle, lead to the non-vanishing contribution. Consider two arbitrary fermions (say, $(k_1\psi)$ and $e(k_2)\psi$) of the vector vertices that couple with some 2 out of 5 fermions of V_5 , say, ψ^{m_1} and ψ^{m_2} . The corresponding factor in the correlator would be given by $k_1^{m_1}e^{m_2}(k_2) + k_1^{m_2}e^{m_1}(k_2)$. But this factor has to be contracted with the tensor B which is antisymmetric in the indices m_1, m_2 . Therefore the total expression vanishes. As for the correlation functions of $V^{(5)}$ with Ramond vertices in the *standard* $-1/2$ -picture one can see that, at least up

to 6-point functions all the correlators vanish (we would like to remind that the Ramond vertex operator in the $-3/2$ -picture is *not* the picture-changed version of the standard picture $-1/2$ fermionic vertex. Rather, it is a different physical state which *does* have non-vanishing matrix elements with V_5 , as it follows from our calculation in the second section of this paper). As for the standard Ramond vertices, all the correlation functions of the type $\langle V^{(-1/2)}(k_1) \dots V^{(-1/2)}(k_{n-1}) V_5 \rangle$ vanish on the sphere at least for $n \leq 6$ due to the ghost anomaly cancelation condition - since the picture-changing factor $f(\Gamma)$ (arising in the measure of the functional integral as a result of integration over supermoduli) which regulates the ghost balance in the correlators, may only increase ghost numbers but not reduce them. For $n > 6$, however, the proof of vanishing of correlation functions of the branelike vertex V_5 with standard Ramond states needs additional explicit computations.

References

- [1] M.Green,J.Schwarz,**Phys.Lett.B136 (1984) 367**
- [2] J.Schwarz**Nucl.Phys.B226(1983) 269**
- [3] R.Metsaev,A.Tseytlin,hep-th/9806095
- [4] R.Kallosh,J.Rahmfeld, A.Rajaraman,hep-th/9805217
- [5] R.Metsaev, A.Tseytlin,**Phys.Lett.B436:281-288,1998**
- [6] R.Kallosh, A.Tseytlin,**J.High Energy Phys.9810:016,1998**
- [7] I.Pesando,**JHEP11(1998)002**
- [8] R.Kallosh, J.Rahmfeld, hep-th/9808038
- [9] S.Gubser,I.Klebanov, A.M.Polyakov, **Phys.Lett.B428:105-114**
- [10] A.M.Polyakov**Nucl.Phys.B486:23-33,1997**
- [11] A.M.Polyakov,hep-th/9809057
- [12] E.Witten**Adv.Theor.Math.Phys.2:253-291,1998**
- [13] J.Maldacena,hep-th/9711200
- [14] D.Polyakov**Phys.Rev.D57:2564-2570,1998**
- [15] J.Polchinski,**Phys.Rev.Lett.75:4724-4727,1995**
- [16] S.Gubser,A.Hashimoto,I.Klebanov,J.Maldacena, **Nucl.Phys.B472:231-248,1996**
- [17] G.Gibbons, P.Townsend**Phys.Rev.Lett.71(1993),5223**